

Jour. Math. Anal.Appl., 258, N1, (2001), 448-456.

Linear ill-posed problems and dynamical systems ^{*†}

A.G. Ramm

Mathematics Department, Kansas State University,
Manhattan, KS 66506-2602, USA
ramm@math.ksu.edu

Abstract

A new approach to solving linear ill-posed problems is proposed. The approach consists of solving a Cauchy problem for a linear operator equation and proving that this problem has a global solution whose limit at infinity solves the original linear equation.

1 Introduction

Let A be a linear, bounded, injective operator on a Hilbert space H , and assume that A^{-1} is unbounded and that $\|A\| \leq \sqrt{m}$, where $m > 0$ is a constant. For example, A may be a compact injective linear operator. Consider the equation,

$$Au = f. \tag{1.1}$$

Assume that (1.1) is solvable, so that $f = Ay$ for a unique $y \in H$. Problem (1.1) is ill-posed since A^{-1} is unbounded. Equation (1.1) cannot be solvable for all $f \in H$ because if A is injective, linear, closed and $R(A) = H$, then A^{-1} must be bounded (by the Banach theorem). Let f_δ be given, such that

$$\|f_\delta - f\| \leq \delta \tag{1.2}$$

^{*}key words: ill-posed problems, dynamical systems, operator equations in Hilbert spaces

[†]Math subject classification: 35R30, 47H17, 65J15

Equation (1.1) with f_δ in place of f may have no solution, and if it has a solution u_δ then it may be that $\|u - u_\delta\|$ is large, although $\delta > 0$ is small. There is a large literature on ill-posed problems since they are important in applications. (See e.g. [4], [3]). In this paper a new approach to solving linear ill-posed problems is proposed. This approach consists of the following steps:

Step 1. Solve the Cauchy problem:

$$\dot{u} = -[Bu + \varepsilon(t)u - F_\delta], \quad u(0) = u_0, \quad (1.3)$$

where

$$\dot{u} := \frac{du}{dt}, \quad B := A^*A, \quad F_\delta := A^*f_\delta, \quad \|F_\delta - F\| \leq \delta\sqrt{m}, \quad F = By,$$

and

$$\varepsilon(t) \in C^1[0, \infty); \quad \varepsilon(t) > 0; \quad \varepsilon(t) \searrow 0 \text{ as } t \rightarrow \infty; \quad \frac{|\dot{\varepsilon}(t)|}{\varepsilon^{\frac{5}{2}}(t)} \rightarrow 0 \text{ as } t \rightarrow \infty. \quad (1.4)$$

One has $\|A^*(f_\delta - f)\| \leq \sqrt{m}\delta$, where we have used the estimate $\|A\| = \|A^*\| \leq \sqrt{m}$.

Examples of functions $\varepsilon(t)$ satisfying (1.4) can be constructed by the formula:

$$\varepsilon(t) = [c + \int_0^t h(s)ds]^{-\frac{2}{3}},$$

where $c > 0$ is a constant, $h(s) > 0$ is a continuous function defined for all $s \geq 0$, such that $h(s) \rightarrow 0$ as $s \rightarrow \infty$ and $\int_0^\infty h(s)ds = \infty$. One has $\frac{|\dot{\varepsilon}(t)|}{\varepsilon^{\frac{5}{2}}(t)} = \frac{2h(t)}{3} \rightarrow 0$ as $t \rightarrow \infty$. For example, $\varepsilon(t) = \frac{1}{\log(t+2)}$ satisfies (1.4). If

$$h(t) = \frac{1}{(2+t)\log(2+t)},$$

then

$$\varepsilon(t) = \frac{1}{(1 + \log \log(2+t))^{\frac{2}{3}}}.$$

This $\varepsilon(t)$ yields nearly fastest decay of $h(t)$ allowed by the restriction $\int_0^\infty h(s)ds = \infty$.

Step 2. Calculate $u(t_\delta)$, where $t_\delta > 0$ is a number which is defined by formula (1.9) below.

Then $t_\delta \rightarrow \infty$ as $\delta \rightarrow 0$ and satisfies the inequality:

$$\|u(t_\delta) - y\| \leq \eta(\delta) \rightarrow 0 \text{ as } \delta \rightarrow 0, \quad (1.5)$$

for a certain function $\eta(\delta) > 0$. If $\delta = 0$, so that $F_\delta = By$, then Step 2 yields the relation

$$\lim_{t \rightarrow \infty} \|u(t) - y\| = 0. \quad (1.6)$$

The foregoing approach is justified in Section 2. Our basic results are formulated as follows.

Theorem 1.1. *Assume that equation (1.1) is uniquely solvable, (1.4) holds, and $\delta = 0$. Then for any u_0 , problem (1.3), with $F = By$ replacing F_δ , has a unique global solution and (1.6) holds.*

By global solution we mean the solution defined for all $t > 0$.

Theorem 1.2. *Assume that equation (1.1) is uniquely solvable, (1.4) holds, and $\delta > 0$. Then for any u_0 problem (1.3) has a unique global solution $u(t)$ and there exists a $t_\delta \rightarrow \infty$ as $\delta \rightarrow 0$, such that $\|u(t_\delta) - y\| \rightarrow 0$ as $\delta \rightarrow 0$. The number t_δ is defined by formula (1.9).*

Let y solve (1.1). Then $By = F := A^*f$ and $\|B\| \leq m$. If

$$\phi(\beta) := \phi(\beta, y) := \beta \left\| \int_0^m \frac{dE_\lambda y}{\lambda + \beta} \right\|, \quad (1.7)$$

where E_λ is the resolution of the identity of the selfadjoint operator B , $E_{\lambda=0} = E_\lambda$, $\beta(\delta)$ is the minimizer of the function

$$h(\beta, \delta) := \phi(\beta) + \frac{\delta}{2\beta^{\frac{1}{2}}} \quad (1.8)$$

on $(0, \infty)$, (see formula (2.20) and Remark 2.3 below), and

$$\eta(\delta) := h(\beta(\delta), \delta), \quad \varepsilon(t_\delta) = \beta(\delta), \quad (1.9)$$

then $t_\delta \rightarrow \infty$ as $\delta \rightarrow 0$, $\eta(\delta) \rightarrow 0$ as $\delta \rightarrow 0$, and

$$\lim_{\delta \rightarrow 0} \|u(t_\delta) - y\| = 0. \quad (1.10)$$

Because B is injective, zero is not an eigenvalue of B , so, for any $y \in H$, one has $\|\int_0^s dE_\lambda y\| \rightarrow 0$ as $s \rightarrow 0$. Therefore $\phi(\beta, y) \rightarrow 0$ as $\beta \rightarrow 0$, for any fixed y . From (2.15) (see below) one gets

$$\|u(t_\delta) - y\| < \eta(\delta) + g_\delta(t_\delta) \rightarrow 0 \text{ as } \delta \rightarrow 0, \quad (1.11)$$

where $g_\delta(t)$ is given by the right-hand side of (2.12) with $\|f_\delta\|$ replacing $\|f\|$.

Remark 1.1. *Theorem 1.2 shows that solving the Cauchy problem (1.3) and calculating its solution at a suitable time t_δ yields a stable solution to ill-posed problem (1.1) and this stable approximate solution satisfies the error estimate (1.11).*

For nonlinear ill-posed problems a similar approach is proposed in [1].

2 Proofs

2.1 Proof of Theorem 1.1

We start with a simple, known fact: if equation (1.1) is solvable, then it is equivalent to the equation

$$Bu = A^*f = By \quad (2.1)$$

Indeed, if $Ay = f$, then apply A^* and get (2.1). Conversely, if (2.1) holds, then $(B(u - y), u - y) = \|A(u - y)\|^2 = 0$, thus $Au = Ay$ and $u = y$, so (1.1) is solvable and its solution is the solution to (2.1). Therefore we will study equation (2.1). The operator $B = A^*A$ is selfadjoint and nonnegative, that is, $(Bu, u) \geq 0$. Let E_λ be its resolution of the identity.

We make another observation: If (1.4) holds, then

$$\int_0^\infty \varepsilon(t) dt = \infty. \quad (2.2)$$

Indeed, (1.4) implies

$$-\frac{\dot{\varepsilon}}{\varepsilon^2} \leq c,$$

where $c = \text{const} > 0$, so

$$\begin{aligned}\frac{d}{dt} \frac{1}{\varepsilon} &\leq c, \\ \frac{1}{\varepsilon(t)} - \frac{1}{\varepsilon(0)} &\leq ct, \\ \frac{1}{\varepsilon(t)} &\leq c_0 + ct, \\ c_0 &:= [\varepsilon(0)]^{-1} > 0,\end{aligned}$$

and

$$\varepsilon(t) \geq \frac{1}{c_0 + ct}.$$

Formula (2.2) follows from the foregoing inequality.

Consider the problem

$$Bw + \varepsilon(t)w - F = 0, \quad F := A^*f = By. \quad (2.3)$$

Since $B \geq 0$ and $\varepsilon(t) > 0$, the solution $w(t)$ of (2.3) exists, is unique and admits the estimate

$$\|w\| \leq \|(B + \varepsilon(t))^{-1}F\| \leq \frac{\|F\|}{\varepsilon(t)}. \quad (2.4)$$

If $F = A^*f$, then (see Remark 2.3 below) one gets:

$$\|w\| \leq \|(B + \varepsilon(t))^{-1}F\| = \|(B + \varepsilon(t))^{-1}A^*f\| \leq \frac{\|f\|}{2\varepsilon^{\frac{1}{2}}(t)}. \quad (2.4')$$

Differentiate (2.3) with respect to t (this is possible by the implicit function theorem) and get

$$[B + \varepsilon(t)]\dot{w} = -\dot{\varepsilon}w, \quad \|\dot{w}\| \leq \frac{|\dot{\varepsilon}|}{\varepsilon}\|w\| \leq \frac{|\dot{\varepsilon}(t)|}{\varepsilon^2(t)}\|F\|, \quad (2.5)$$

where (2.4) was used.

Using (2.4') yields:

$$\|\dot{w}\| \leq \frac{|\dot{\varepsilon}|}{\varepsilon}\|w\| \leq \frac{|\dot{\varepsilon}(t)|}{2\varepsilon^{\frac{3}{2}}(t)}\|f\|. \quad (2.5')$$

Denote

$$z(t) := u(t) - w(t). \quad (2.6)$$

Subtract (2.3) from (1.3) (with F in place of F_δ) and get

$$\dot{z} = -\dot{w} - [B + \varepsilon(t)]z, \quad z(0) = u_0 - w(0). \quad (2.7)$$

Multiply (2.7) by $z(t)$ and get

$$(\dot{z}, z) = -(\dot{w}, z) - (Bz, z) - \varepsilon(t)(z, z). \quad (2.8)$$

Denote

$$\|z(t)\| := g(t) \quad (2.9)$$

Then the inequality $(Bz, z) \geq 0$ and equation (2.8) imply:

$$g\dot{g} \leq \|\dot{w}\|g - \varepsilon(t)g^2. \quad (2.10)$$

Because $g \geq 0$, it follows from (2.10) and (2.5') that

$$\dot{g} \leq \|f\| \frac{|\dot{\varepsilon}(t)|}{2\varepsilon^{\frac{3}{2}}(t)} - \varepsilon(t)g(t), \quad g(0) = \|u_0 - w_0\|, \quad (2.11)$$

so

$$g(t) \leq e^{-\int_0^t \varepsilon(s)ds} \left[g(0) + \int_0^t e^{\int_0^\tau \varepsilon(s)ds} \frac{|\dot{\varepsilon}(\tau)|}{2\varepsilon^{\frac{3}{2}}(\tau)} d\tau \|f\| \right]. \quad (2.12)$$

Assumption (1.4) (the last one in (1.4)) and (2.12) imply (use L'Hospital's rule) that

$$\|u(t) - w(t)\| := g(t) \rightarrow 0 \text{ as } t \rightarrow +\infty. \quad (2.13)$$

The existence of the global solution to (1.3) is obvious since equation (1.3) is linear and the operator B is bounded.

To prove (1.6) it is sufficient to prove that

$$\|w(t) - y\| \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (2.14)$$

Indeed, if (2.14) holds then (2.13) and (2.14) imply:

$$\|u(t) - y\| \leq \|u(t) - w(t)\| + \|w(t) - y\| \rightarrow 0 \text{ as } t \rightarrow \infty. \quad (2.15)$$

We now prove (2.14). One has:

$$\|w(t) - y\| = \left\| \int_0^m \frac{\lambda}{\lambda + \varepsilon(t)} dE_\lambda y - \int_0^m dE_\lambda y \right\| = \left\| \int_0^m \frac{\varepsilon(t)}{\lambda + \varepsilon(t)} dE_\lambda y \right\|. \quad (2.16)$$

Thus

$$\|w(t) - y\| = \phi(\varepsilon(t), y), \quad (2.17)$$

where $\phi(\varepsilon, y) := \phi(\varepsilon)$ is as defined in (1.7). Since B is injective, the point $\lambda = 0$ is not an eigenvalue of B . Therefore

$$\lim_{\varepsilon \rightarrow 0} \phi(\varepsilon) = 0, \quad (2.18)$$

by the Lebesgue dominant convergence theorem.

Thus (2.14) follows and Theorem 1.1 is proved. $\square$

2.2 Proof of Theorem 1.2

The proof is quite similar to the above, so we indicate only the new points. Equation (2.3) is now replaced by the equation

$$Bw + \varepsilon(t)w - F_\delta = 0. \quad (2.19)$$

Estimates (2.4), (2.4'), (2.5), (2.5') and (2.13) hold with F_δ and f_δ in place of F and f , respectively. The main new point is the estimate of $w(t) - y$:

$$\begin{aligned} \|w(t) - y\| &= \left\| \int_0^m \frac{dE_\lambda F_\delta}{\lambda + \varepsilon(t)} - \int_0^m dE_\lambda y \right\| \\ &= \left\| \int_0^m \frac{dE_\lambda (F_\delta - F)}{\lambda + \varepsilon(t)} \right\| + \phi(\varepsilon(t)) \\ &\leq \phi(\varepsilon(t)) + \frac{\delta}{2\varepsilon^{\frac{1}{2}}(t)}, \end{aligned} \quad (2.20)$$

where $\|f - f_\delta\| \leq \delta$ and estimate (2.4') was used.

If $\beta(\delta)$ is the minimizer of the function (1.8), then

$$h(\beta(\delta), \delta) := \eta(\delta) \rightarrow 0 \text{ as } \delta \rightarrow 0; \quad \beta(\delta) \rightarrow 0 \text{ as } \delta \rightarrow 0. \quad (2.21)$$

The latter relation in (2.21) holds because $\phi(\beta) \rightarrow 0$ as $\beta \rightarrow 0$.

Since $\varepsilon(t) \searrow 0$ as $t \rightarrow \infty$, one can find the unique t_δ such that

$$\varepsilon(t_\delta) = \beta(\delta) \rightarrow 0 \text{ as } \delta \rightarrow 0. \quad (2.22)$$

Thus

$$\|w(t_\delta) - y\| \leq \eta(\delta) \rightarrow 0 \text{ as } \delta \rightarrow 0. \quad (2.23)$$

The function $\eta(\delta) = \eta(\delta, y)$ depends on y because $\phi(\varepsilon) = \phi(\varepsilon, y)$ does (see formula (1.7)).

Combining (2.23), (2.13) and (2.15) one gets the conclusion of Theorem 1.2. $\square$

Remark 2.1. *We also give a proof of (2.14) which does not use the spectral theorem.*

From (2.3) one gets

$$Bx + \varepsilon(t)x = -\varepsilon(t)y, \quad x(t) := w(t) - y. \quad (2.24)$$

Thus $(Bx, x) + \varepsilon(x, x) = -\varepsilon(y, x)$. Since $(Bx, x) \geq 0$ and $\varepsilon > 0$, one gets

$$(x, x) \leq |(y, x)|, \quad \|x(t)\| \leq \|y\| = \text{const} < \infty. \quad (2.25)$$

Bounded sets in H are weakly compact. Therefore there exists a sequence $t_n \rightarrow \infty$ such that

$$x_n := x(t_n) \rightharpoonup x_\infty, \quad n \rightarrow \infty \quad (2.26)$$

where $\rightharpoonup$ stands for the weak convergence. From (2.24) and (2.25) it follows that

$$Bx_n \rightarrow 0, \quad n \rightarrow \infty. \quad (2.27)$$

A monotone hemicontinuous operator is weakly closed. This claim, which we prove below, implies that (2.26) and (2.27) yield $Bx_\infty = 0$. because B is injective, $x_\infty = 0$, that is, $x(t_n) \rightharpoonup 0$. From (2.25) it follows that $\|x(t_n)\| \rightarrow 0$

as $n \rightarrow \infty$, because $(y, x(t_n)) \rightarrow 0$ as $n \rightarrow \infty$, due to $x(t_n) \rightarrow 0$. By the uniqueness of the limit, one concludes that $\lim_{t \rightarrow \infty} \|x(t)\| = 0$, which is (2.14).

Let us now prove the claim.

We wish to prove that $x_n \rightharpoonup x$ and $Bx_n \rightarrow f$ imply $Bx = f$ provided that B is monotone and hemicontinuous. The monotonicity implies $(Bx_n - B(x - \varepsilon p), x_n - x + \varepsilon p) \geq 0$ for all $\varepsilon > 0$ and all $p \in H$. Take $\varepsilon \rightarrow 0$ and use hemicontinuity of B to get $(f - Bx, p) \geq 0 \quad \forall p \in H$. Take $p = f - Bx$ to obtain $Bx = f$, as claimed. $\square$

The above argument uses standard properties of monotone hemicontinuous operators [2].

Remark 2.2. In (2.23) $\eta(\delta) = O(\delta^{\frac{2}{3}})$ is independent of y if y runs through a set $S_a := S_{a,R} := \{y : y = B^a h, \|h\| \leq R\}$, where $R > 0$ is an arbitrary large fixed number and $a \geq 1$. If $0 < a < 1$, then $\eta(\delta) = O(\delta^{\frac{2a}{2a+1}})$, as $\delta \rightarrow 0$, and this estimate is uniform with respect to $y \in S_{a,R}$.

Consider, for example, the case $a \geq 1$. If $y = B^a h$, then $\phi(\varepsilon)$ in (2.20) can be chosen for all $y \in S_{a,R}$ simultaneously. Using (1.7), one gets:

$$\phi(\varepsilon) = \varepsilon \sup_{\|h\| \leq R} \left\| \int_0^m \frac{\lambda^a}{\lambda + \varepsilon} dE_\lambda h \right\| \leq \varepsilon m^{a-1} R,$$

where $a \geq 1$ and ε is positive and small. For a fixed $\delta > 0$ one finds the minimizer $\varepsilon(\delta) = O(\delta^{\frac{2}{3}})$ of the function $\frac{\delta}{2\varepsilon^{\frac{1}{2}}} + \varepsilon m^{a-1} R$ and the minimal value $\eta(\delta)$ of this function is $O(\delta^{\frac{2}{3}})$.

If B is compact, then the condition $y \in S_a$ means that y belongs to a compactum which is the image of a bounded set $\|h\| \leq R$ under the map B^a .

The case $0 < a < 1$ is left to the reader. It can be treated by the method used above.

Remark 2.3. It can be checked easily that

$$A(A^*A + \epsilon I)^{-1} = (AA^* + \epsilon I)^{-1}A.$$

This implies

$$\|(B + \epsilon I)^{-1}A^*f\|^2 = ((b + \epsilon I)^{-2}bf, f) := J,$$

where $B := A^*A$ and $b := AA^* \geq 0$.

Thus,

$$J = \int_0^m s(s + \epsilon)^{-2} d(e_s f, f) \leq \frac{1}{4\epsilon} \|f\|^2,$$

where e_s is the resolution of the identity corresponding to the selfadjoint operator b . Therefore one gets the following estimate:

$$\|(B + \epsilon I)^{-1} A^* f\| \leq \frac{1}{2\sqrt{\epsilon}} \|f\|.$$

This estimate was used to obtain estimates (2.4') and (2.5'). For example, estimate (2.4) was replaced by the following one:

$$\|(B + \epsilon I)^{-1} F\| \leq \frac{1}{2\epsilon^{\frac{1}{2}}} \|f\|, \quad (2.4')$$

and (2.5) can be replaced by the estimate:

$$\|\dot{w}\| \leq \frac{|\dot{\epsilon}|}{\epsilon} \|w\| \leq \frac{|\dot{\epsilon}(t)|}{2\epsilon^{3/2}(t)} \|f\|. \quad (2.5')$$

These estimates were used to improve the estimate for $\eta(\delta)$ in the previous remark.

Estimate (2.4') was used by a suggestion of a referee. The author thanks the referee for the suggestion.

In fact, one can prove a stronger estimate than (2.4'), namely $\|w\| \leq \|y\|$. Indeed, multiply (2.3) by $w - y$, use the nonnegativity of B and positivity of ϵ and get $(w, w - y) \leq 0$. Thus $\|w\|^2 \leq \|w\| \|y\|$, and the desired inequality $\|w\| \leq \|y\|$ follows.

Appendix. Let us give an alternative proof of Theorem 1.2. Let $u_\delta(t)$ solve (1.3), $u(t)$ solve (1.3) with F_δ replaced by F , and $u_\delta(t)$ and $u(t)$ satisfy the same initial condition. Denote $w_\delta := u_\delta(t) - u(t)$ and let $\|w_\delta\| := g_\delta(t)$. One has:

$$\dot{w}_\delta = -[Bw_\delta + \epsilon(t)w_\delta - h_\delta], \quad w_\delta(0) = 0,$$

where $h_\delta := F_\delta - F$, $\|h_\delta\| < \sqrt{m}\delta := c\delta$. Multiply the above equation by w_δ in H , use the inequality $B \geq 0$ and get

$$\dot{g}_\delta \leq -\epsilon(t)g_\delta + c\delta.$$

Since $g_\delta(0) = 0$, this implies:

$$g_\delta(t) \leq c\delta \exp\left[-\int_0^t \epsilon(s)ds\right] \int_0^t \exp\left[\int_0^p \epsilon(s)ds\right]dp \leq c\frac{\delta}{\epsilon(t)}.$$

Thus

$$\|u_\delta(t) - y\| \leq \|u_\delta(t) - u(t)\| + \|u(t) - y\| \leq c\frac{\delta}{\epsilon(t)} + a(t),$$

where $a(t) := \|u(t) - y\| \rightarrow 0$ as $t \rightarrow \infty$. Define t_δ as the minimal minimizer of the following function of t for a fixed $\delta > 0$:

$$c\frac{\delta}{\epsilon(t)} + a(t) = \min := \mu(\delta).$$

Since $a(t) \rightarrow 0$ and $\epsilon(t) \rightarrow 0$ as $t \rightarrow \infty$, one concludes that the minimal minimizer $t_\delta \rightarrow \infty$ as $\delta \rightarrow 0$ and $\mu(\delta) \rightarrow 0$ as $\delta \rightarrow 0$. Theorem 1.2 is proved $\square$.

References

- [1] Airapetyan, R., Ramm, A.G., Dynamical systems and discrete methods for solving nonlinear ill-posed problems, Appl.Math.Rev.,Vol.1, World Sci. Publishers, (ed. G.Anastassiou), (2000), pp.491-536.
- [2] Deimling, K., **Nonlinear functional analysis**, Springer Verlag, Berlin, 1985
- [3] Engl, W., Hanke, M., Neubauer, A., **Regularization of ill-posed problems**, Kluwer, Dordrecht, 1996.
- [4] Groetsch, C., **Inverse problems in mathematical sciences**, Vieweg, Braunschweig, 1993.